产品名称：Recombinant Human CD200 R1 (C-Fc)
产品货号：PHH0308

概述（Summary）

英文全称	CD200 R1
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	Recombinant Human CD200 Receptor 1 is produced by our Mammalian expression system and the target gene encoding Ala27-Leu266 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q8TD46
表达宿主（Host）	Human Cells
种属（Species）	Human
预测分子量（Predicted MW）	53.9 KDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
运输方式（Shipping）	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图（SDS-PAGE image）

产品名称：Recombinant Human CD200 R1 (C-Fc)
产品货号：PHH0308



背景（Background）

分子别名（Alternative Names）

Cell Surface Glycoprotein CD200 Receptor 1; CD200 Cell Surface Glycoprotein Receptor; Cell Surface Glycoprotein OX2 Receptor 1; CD200R1; CD200R; CRTR2; MOX2R; OX2R

背景介绍（References）

Cell surface glycoprotein CD200 Receptor 1 (CD200R1) is the receptor for the CD200 (OX-2) membrane glycoprotein. CD200R1 contains one C2- type Ig-like domain and one V-type Ig-like domain within its extracellular domain and a PTB-signaling motif in cytoplasmic domain. CD200R1 and CD200 associate via their respective N-terminal Ig-like domains. CD200R1 is restricted primarily to mast cells, basophils, macrophages, and dendritic cells. It propagates inhibitory signals despite its lacking a cytoplasmic ITIM (immunoreceptor tyrosinebased inhibitory motif). The receptor-substrate interaction may function as a myeloid downregulatory signal.

注意事项（Note）

For research use only .