


Product Name: Recombinant Human CD26 (N-Fc)
Catalog #: PHH0316



概述 (Summary)

英文全称	DPPIV/CD26/DPP4/Dipeptidyl Peptidase 4
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Dipeptidyl Peptidase 4 is produced by our Mammalian expression system and the target gene encoding Asn29-Pro766 is expressed with a human IgG1 Fc tag at the N-terminus.
Accession #	P27487
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	111.7 KDa
蛋白形态 (Form)	Supplied as a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	

电泳图 (SDS-PAGE image)



背景 (Background)

分子别名 (Alternative Names) Dipeptidyl peptidase 4; ADABP; Adenosine deaminase complexing protein 2;

Product Name: Recombinant Human CD26 (N-Fc)
Catalog #: PHH0316



背景介绍 (References)

ADCP-2; Dipeptidyl peptidase IV; DPP IV; T-cell activation antigen CD26

CD26 is a signal-anchor for type II membrane protein that belongs to the peptidase S9B family. CD26 is expressed specifically in lymphatic vessels but not in blood vessels in the skin, small intestine, esophagus, ovary, breast and prostate glands. It acts as a positive regulator of T-cell coactivation, by binding at least ADA, CAV1, IGF2R, and PTPRC. It' s binding to CAV1 and CARD11 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner. Its interaction with ADA also regulates lymphocyte-epithelial cell adhesion. In association with FAP is involved in the pericellular proteolysis of the extracellular matrix (ECM), the migration and invasion of endothelial cells into the ECM. It may be involved in the promotion of lymphatic endothelial cells adhesion, migration and tube formation. When overexpressed, it enhanced cell proliferation, a process inhibited by GPC3. It acts also as a serine exopeptidase with a dipeptidyl peptidase activity that regulates various physiological processes by cleaving peptides in the circulation, including many chemokines, mitogenic growth factors, neuropeptides and peptide hormones.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.