Catalog #: PHH0326



概述 (Summary)

CLEC10A/CD301 英文全称

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/µg as determined by LAL test.

蛋白构建 (Construction) Recombinant Human C-Type Lectin Domain Family 10 Member A is

produced by our Mammalian expression system and the target gene

encoding Gln61-His316 is expressed with a 6His tag at the C-terminus.

Accession # Q8IUN9

蛋白标签 (Tag)

表达宿主 (Host) **Human Cells**

种属 (Species) Human 预测分子量 (Predicted MW) 29.8 KDa

Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. 蛋白形态 (Form)

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 稳定性&储存 (Stability &Storage)

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

> is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized

> protein in distilled water. Please aliquot the reconstituted solution to minimize


freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human CLEC10A (C-6His) Catalog #: PHH0326





背景 (Background)

分子别名 (Alternative Names)

背景介绍 (References)

C-Type Lectin Domain Family 10 Member A; C-Type Lectin Superfamily Member 14; Macrophage Lectin 2; CD301; CLEC10A; CLECSF13; CLECSF14; HML

C-Type Lectin Domain Family 10 Member A (CLEC10A) is a type II transmembrane C-type lectin that is expressed on immature myleloid dendritic cells and alternatively activated (tolerogenic) macrophages. CLEC10A/MGL binds and internalizes molecules with terminal nonsialylated GalNAc carbohydrates such as the Tn carcinoma antigen. CLEC10A/MGL also binds the GP envelope glycoprotein on Marburg and Ebola viruses and enhances viral entry and infectivity. It constitute a unique class of C-type lectins because of their specificity for galactose and its structural homologues. CLEC10A is thought to participate in the recognition of molecules from both altered self and pathogens due to its monosaccharide specificity for Gal and N-acetylgalactosamine (GalNAc). Human and rat carry a single gene for CLEC10A/MGL, while mouse has two closely related MGL1 and MGL2 genes.

注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.