


概述 (Summary)

英文全称	CD320/TCbIR/8D6A
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Transcobalamin II Receptor is produced by our Mammalian expression system and the target gene encoding Ser36-Val231 is expressed with a 6His tag at the C-terminus.
Accession #	Q9NPF0
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	21.1 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 10mM Tris-Citrate, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human TCbIR (C-6His)
Catalog #: PHH0332



背景 (Background)

分子别名 (Alternative Names)

CD320 antigen;8D6 antigen;FDC-signaling molecule 8D6;FDC-SM-8D6;Transcobalamin receptor;TCbIR;CD320

背景介绍 (References)

CD320 antigen is also known as 8D6 antigen , FDC-signaling molecule 8D6 , Transcobalamin receptor and 8D6A. It is a single-pass type I membrane protein and containing two LDL-receptor class A domains. CD320 has been recently discovered and reported as a follicular dendritic cell (FDC) protein. CD320 can augments the proliferation of plasma cells precursors generated by IL-10. CD320 also founctions a receptor for the cellular uptake of transcobalamin bound cobalamin. Defects in CD320 are the cause of methylmalonic aciduria type TCbIR (MMATC) which is a metabolic disorder.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.