


概述 (Summary)

英文全称	CD47/IAP/OA3/Leukocyte Surface Antigen CD47/Antigenic surface determinant protein OA3/Integrin-associated protein/Protein MER6
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Leukocyte Surface Antigen CD47 is produced by our Mammalian expression system and the target gene encoding Gln19-Pro139 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q08722
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	40.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human CD47 (C-Fc)
Catalog #: PHH0354



背景 (Background)

分子别名 (Alternative Names)

Leukocyte Surface Antigen CD47; Antigenic Surface Determinant Protein OA3; Integrin-Associated Protein; IAP; Protein MER6; CD47; MER6

背景介绍 (References)

CD47(Integrin-Associated Protein,IAP) is a 40 - 60 kDa variably glycosylated atypical member of the immunoglobulin superfamily. The ubiquitously expressed CD47 binds to SIRP family members on macrophages, neutrophils, and T cells. CD47 is involved in the increase in intracellular calcium concentration that occurs upon cell adhesion to extracellular matrix. The protein is also a receptor for the C-terminal cell-binding domain of thrombospondin, and it may play a role in membrane transport and signal transduction. This protein has broad tissue distribution, and is reduced in expression on Rh erythrocytes.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.