


## 概述 (Summary)

|                             |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|-----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 英文全称                        | CD55/DAF                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| 纯度 (Purity)                 | Greater than 95% as determined by reducing SDS-PAGE                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| 内毒素 (Endotoxin level)       | <1 EU/μg as determined by LAL test.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| 蛋白构建 (Construction)         | Recombinant Human CD55 is produced by our Mammalian expression system and the target gene encoding Asp35-Ser353 is expressed with a 6His tag at the C-terminus.                                                                                                                                                                                                                                                                                                                                                                                                               |
| Accession #                 | P08174                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| 蛋白标签 (Tag)                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| 表达宿主 (Host)                 | Human Cells                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| 种属 (Species)                | Human                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| 预测分子量 (Predicted MW)        | 36 KDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| 蛋白形态 (Form)                 | Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| 储存缓冲液 (Buffer)              |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| 运输方式 (Shipping)             | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| 稳定性&储存 (Stability &Storage) | Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.                                                                                                                                                                                                                                                                                                                                                                                                          |
| 复溶 (Reconstitution)         | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

## 电泳图 (SDS-PAGE image)

**Product Name: Recombinant Human CD55 (C-6His)**  
**Catalog #: PHH0359**



## 背景 (Background)

### 分子别名 (Alternative Names)

Complement Decay-Accelerating factor; CD55; CR; DAF

### 背景介绍 (References)

CD55 is a member of the RCA (regulators of complement activation) family. RCA proteins is characterized by the presence of four to 30 SCRs (short consensus repeats also called CCPs for control protein modules) in their plasmaexposed regions. CD55 containing four SCR modules is involved in the regulation of the complement cascade. CD55 is known to bind CD97 via the first SCR. It also binds physiologically generated C3 convertases with its second and third SCRs. Binding results in an accelerated "decay" , or dissociation of active C3 convertases, thus blocking the development of C' attack complexes on nonforeign cells. It is known that viruses and bacteria also utilize multiple SCR sites for infection.

## 注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.