


概述 (Summary)

英文全称	CHRNB3/Neuronal acetylcholine receptor subunit beta-3
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Neuronal Acetylcholine Receptor Subunit Beta-3 is produced by our Mammalian expression system and the target gene encoding Ile25-Leu232 is expressed with a 6His tag at the C-terminus.
Accession #	Q05901
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	25.3 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human CHRNB3 (C-6His)
Catalog #: PHH0408



背景 (Background)

分子别名 (Alternative Names)

Neuronal acetylcholine receptor subunit beta-3

背景介绍 (References)

Neuronal acetylcholine receptor subunit beta-3(CHRNB3) is a cell membrane protein and belongs to the ligand-gated ion channel (TC 1.A.9) family. CHRNB3 seems to be composed of two different type of subunits: alpha and beta. The CHRNB3 are (hetero) pentamers composed of homologous subunits. The subunits that make up the muscle and neuronal forms of CHRNB3 are encoded by separate genes and have different primary structure. There are several subtypes of neuronal CHRNB3 that vary based on which homologous subunits are arranged around the central channel. They are classified as alpha-subunits if like muscle alpha-1, they have a pair of adjacent cysteines as part of the presumed acetylcholine binding site. Subunits lacking these cysteine residues are classified as beta-subunits.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.