

Product Name: Recombinant Human Cryptic (C-6His)
Catalog #: PHH0455




概述 (Summary)

英文全称	Cryptic protein/cryptic
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Cryptic Protein is produced by our Mammalian expression system and the target gene encoding Tyr26-Gly169 is expressed with a 6His tag at the C-terminus.
Accession #	P0CG37
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	16.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human Cryptic (C-6His)
Catalog #: PHH0455



背景 (Background)

分子别名 (Alternative Names)

Cryptic protein;Cryptic family protein 1;CFC1

背景介绍 (References)

Cryptic (CFC1) is a member of the epidermal growth factor (EGF)- Cripto, Frl-1, and Cryptic (CFC) family. It contains an EGF-like domain, and is glycosylated on its N-terminal during post-translational modification. Cryptic is identified as a NODAL coreceptor involved in the correct establishment of the left-right axis. It may play a role in mesoderm and/or neural patterning during gastrulation. The unnormal expression of this gene may causes a series of diseases such as HTX2, Transposition of the great arteries dextro-looped 2, and Conotruncal heart malformations.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.