


概述 (Summary)

英文全称	CXCL8/IL-8 (Glu21-Ser99)
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human C-X-C Motif Chemokine 8/Interleukin 8 is produced by our Mammalian expression system and the target gene encoding Glu21-Ser99 is expressed with a 6His tag at the C-terminus.
Accession #	P10145
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	10.1 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)



背景 (Background)

分子別名 (Alternative Names)

Interleukin-8; IL-8; C-X-C Motif Chemokine 8; CXCL8; Emoctakin; Granulocyte Chemotactic Protein 1; GCP-1; Monocyte-Derived Neutrophil Chemotactic Factor; MDNCF; Monocyte-Derived Neutrophil-Activating Peptide; MONAP; Neutrophil-Activating Protein 1; NAP-1

背景介绍 (References)

Interleukin-8 (IL-8) belongs to the neutrophil-specific CXC family of chemokines. It is one of the initial cytokines released from a variety of cell types, including T cells, endothelial cells and fibroblasts, in response to an inflammatory stimulus and acts by recruiting neutrophils, T-cells and basophils to the site of inflammation. Elevated Interleukin-8 levels are associated with the onset of a variety of disease states.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.