Product Name: Recombinant Human Decorin (C-6His)

Catalog #: PHH0522



概述 (Summary)

英文全称 Decorin

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/μg as determined by LAL test.

蛋白构建 (Construction) Recombinant Human Decorin is produced by our Mammalian expression

system and the target gene encoding Gly17-Lys359 is expressed with a

6His tag at the C-terminus.

Accession # P07585

蛋白标签 (Tag)

表达宿主 (Host) Human Cells

种属 (Species)Human预测分子量 (Predicted MW)38.8 KDa

蛋白形态 (Form) Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

稳定性&储存 (Stability &Storage) Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized

protein in distilled water. Please aliquot the reconstituted solution to minimize

freeze-thaw cycles.


电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human Decorin (C-6His)

Catalog #: PHH0522





背景 (Background)

分子別名 (Alternative Names) 背景介绍 (References) Decorin; Bone proteoglycan II; PG-S2; PG40; DCN; SLRR1B

Decorin is a secreted chondroitin/dermatan sulfate proteoglycan in the family of small leucine-rich proteoglycans (SLRPs). SLRP family members are characterized by N-terminal and C-terminal cysteine-rich regions which flank the central region containing 10-12 tandem leucine-rich repeats (LRR). The human Decorin cDNA encodes a 359 amino acid (aa) precursor that includes a 16 aa signal sequence and a 14 aa propeptide. Alternate splicing of human Decorin generates five isoforms with variable length deletions. Decorin is an N-glycosylated protein that also carries a variablysized hybrid chondroitin/dermatan sulfate chain at Ser34. Decorin regulates assembly of the extracellular collagen matrix and the bioactivity of the matrix associated growth factors FGF2,GDF8/Myostatin, TGFβ,and WISP1. It also binds and activates EGF R, ErbB4, and IGFI-R. In vivo, Decorin promotes myoblast differentiation, supports angiogenesis, and inhibits tumor progression.

注意事项(Note)

For Research Use Only, Not for Diagnostic Use.