Product Name: Recombinant Human DPT (C-Fc-6His)

Catalog #: PHH0529



概述 (Summary)

英文全称 Dermatopontin/DPT

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/μg as determined by LAL test.

蛋白构建 (Construction) Recombinant Human Dermatopontin is produced by our Mammalian

expression system and the target gene encoding Gln19-Val201 is

expressed with a human IgG1 Fc, 6His tag at the C-terminus.

Accession # Q07507

蛋白标签 (Tag)

表达宿主 (Host) Human Cells

种属 (Species) Human 预测分子量 (Predicted MW) 49.9 KDa

蛋白形态 (Form) Lyophilized from a 0.2 µm filtered solution of 20mM PB, 4% Sucrose, 4%

mannitol, 0.02% Tween80, pH 7.5.

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

稳定性&储存 (Stability &Storage) Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized

is not recommended to reconstitute to a concentration less than 100µg/ml.

protein in distilled water. Please aliquot the reconstituted solution to minimize

freeze-thaw cycles.


电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human DPT (C-Fc-6His)

Catalog #: PHH0529





背景 (Background)

分子別名 (Alternative Names) 背景介绍 (References) Dermatopontin; Tyrosine-rich acidic matrix protein; TRAMP and DPT Dermatopontin, also known as Tyrosine-rich acidic matrix protein, TRAMP and DPT, is a secreted protein which belongs to the dermatopontin family. DPT is expressed in various tissues, such as fibroblasts, heart, skeletal muscle, brain and pancreas. It seems to mediate adhesion by cell surface integrin binding. DPT may serve as a communication link between the dermal fibroblast cell surface and its extracellular matrix environment. DPT can enhance TGFB1 activity through interaction with decorin. In addition, DPT accelerates collagen fibril formation, stabilizes collagen fibrils against low-temperature dissociation and inhibits cell proliferation.

注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.