


概述 (Summary)

英文全称	EphA4/Ephrin type-A receptor 4/HEK8/SEK/TYRO1
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Ephrin Type A Receptor 4 is produced by our Mammalian expression system and the target gene encoding Val20-Thr547 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	P54764
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	85.6 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human EphA4 (C-Fc)
Catalog #: PHH0579



背景 (Background)

分子别名 (Alternative Names)

Ephrin type-A receptor 4; HEK8; SEK; TYRO1; EPHA4; Tyrosine-protein kinase receptor SEK; Tyrosine-protein kinase TYRO1; EK8; hEK8; EPH-like kinase 8

背景介绍 (References)

Ephrin type-A receptor 4 (EPHA4) belongs to the protein kinase superfamily and Ephrin receptor subfamily. EPHA4 contains 1 Eph LBD domain, 2 fibronectin type-III domains, 1 protein kinase domain and 1 SAM domain. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.