


概述 (Summary)

英文全称	Ephrin-B2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Ephrin-B2 is produced by our Mammalian expression system and the target gene encoding Ile28-Ala229 is expressed with a 6His tag at the C-terminus.
Accession #	P52799
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	23.24 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human EFNB2 (C-6His)
Catalog #: PHH0596



背景 (Background)

分子别名 (Alternative Names)

Ephrin-B2; EPH-Related Receptor Tyrosine Kinase Ligand 5; LERK-5; HTK Ligand; HTK-L; EFNB2; EPLG5; HTKL; LERK5

背景介绍 (References)

Ephrin-B2 is a type I transmembrane protein and belongs the Ephrin family. It binds to the receptor tyrosine kinases, such as EPHA4, EPHB4 and EPHA3. Ephrin-B2 has been implicated in mediating developmental events, especially in the nervous system, erythropoiesis and tumour metastasis. Ligation of Ephrin-B2 with complementary EphB receptors on adjacent cells results in a combination of forward (EphB receptors) and reverse (Ephrin-B2) signalling, which is central to tissue development and remodelling functions. In addition, Ephrin-B2 may play a role in constraining the orientation of longitudinally projecting axons.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.