Product Name: Recombinant Human EGFR vIII (C-Fc)

Catalog #: PHH0599



概述 (Summary)

英文全称 EGFRvIII

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/μg as determined by LAL test.

蛋白构建 (Construction) Recombinant Human Epidermal Growth Factor Receptor/Receptor

Tyrosine Protein Kinase ErbB1 is produced by our Mammalian expression system and the target gene encoding Leu25-Ser378 is expressed with a

human IgG1 Fc tag at the C-terminus.

Accession # NP 001333870

蛋白标签 (Tag)

表达宿主 (Host) Human Cells 种属 (Species) Human

预测分子量 (Predicted MW) 65.8 KDa

蛋白形态 (Form) Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

稳定性&储存 (Stability &Storage) Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to

is not recommended to reconstitute to a concentration less than 100µg/ml.

reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized

protein in distilled water. Please aliquot the reconstituted solution to minimize


freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human EGFR vIII (C-Fc) Catalog #: PHH0599

C EnkiLife



背景 (Background)

分子别名 (Alternative Names)

背景介绍 (References)

Epidermal growth factor receptor; EGFR; Proto-oncogene c-ErbB-1; Receptor tyrosine-protein kinase erbB-1; EGFR

The EGFR subfamily of receptor tyrosine kinases is composed of EGFR, ErbB2, ErbB3 and ErbB4. The EGFR shares 43% - 44% aa sequence identity with the ECD of human EGFR subfamily. All these family members are type I transmembrane glycoproteins with an extracellular ligand binding domain. The extracellular ligand binding domain is containing two cysteine-rich domains separated by a spacer region and a cytoplasmic domain containing a membrane-proximal tyrosine kinase domain. Ligand binding could induce EGFR homodimerization and heterodimerization with ErbB2, resulting in cell signaling, heterodimerization tyrosine phosphorylation and kinase activation. It can bind EGF, amphiregulin, TGF-alpha, betacellulin, epiregulin, HB-EGF, epigen, and so on. Its signaling regulates multiple biological functions including cell proliferation, differentiation, motility, and apoptosis. EGFR can also be recruited to form heterodimers with the ligand-activated ErbB3 or ErbB4. EGFR is overexpressed in different tumors. Several anti-cancer drugs use EGFR as target.

注意事项(Note)

For Research Use Only, Not for Diagnostic Use.