


Product Name: Recombinant Human GALNTL1 (C-6His)
Catalog #: PHH0713



概述 (Summary)

英文全称	GALNTL1/GalNAc-T-like protein 1
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Putative Polypeptide N-Acetylgalactosaminyltransferase-Like Protein 1 is produced by our Mammalian expression system and the target gene encoding Asp27-Thr558 is expressed with a 6His tag at the C-terminus.
Accession #	Q8N428
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	61 KDa
蛋白形态 (Form)	Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 7.5.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	

电泳图 (SDS-PAGE image)



背景 (Background)

Product Name: Recombinant Human GALNTL1 (C-6His)
Catalog #: PHH0713



分子别名 (Alternative Names)

Putative Polypeptide N-Acetylgalactosaminyltransferase-Like Protein 1; Polypeptide GalNAc Transferase-Like Protein 1; GalNAc-T-Like Protein 1; pp-GaNTase-Like Protein 1; Protein-UDP Acetylgalactosaminyltransferase-Like Protein 1; UDP-GalNAc:Polypeptide N-Acetylgalactosaminyltransferase-Like Protein 1; GALNTL1; KIAA1130

背景介绍 (References)

Putative polypeptide N-acetylgalactosaminyltransferase-like protein 1, also known as Polypeptide GalNAc transferase-like protein 1, Protein-UDP acetylgalactosaminyltransferase-like protein 1, UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase-like protein 1, GalNAc-T-like protein 1, pp-GaNTase-like protein 1 and GALNTL1, belongs to the glycosyltransferase 2 family. GALNTL1 plays an important role in the protein modification and protein glycosylation process. GALNTL1 uses the manganese and calcium ion as a cofactor, may catalyze the initial reaction in O-linked oligosaccharide biosynthesis, transfers the N-acetyl-D-galactosamine residue to a serine or threonine residue on the protein receptor.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.