


概述 (Summary)

英文全称	Glycoprotein A33/GPA33
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Glycoprotein A33 is produced by our Mammalian expression system and the target gene encoding Ile22-Val235 is expressed with a 6His tag at the C-terminus.
Accession #	Q99795
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	24.66 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20 mM PB, 150 mM NaCl, 1mM EDTA, pH7.2.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human GPA33 (C-6His)
Catalog #: PHH0744



背景 (Background)

分子别名 (Alternative Names)

Cell Surface A33 Antigen; Glycoprotein A33; GPA33

背景介绍 (References)

Human Glycoprotein A33 (GPA33) is a single-pass type I membrane protein, belongs to the CTX family of cell adhesion molecular within the immunoglobulin family, can be expressed in normal gastrointestinal epithelium and in 95% of colon cancers. GPA33 consists of one Ig-like C2-type domain and one Ig-like V-type domain. The predicted mature protein includes a single transmembrane domain, a extracellular region and a intracellular tail. Intracellular traffic and recycling to the cell surface appear to play an important role in GPA33 function and to have an influence on its surface density superseding translation regulation. GPA33 has become a promising target of immunologic therapy strategies. GPA33 may also play a important role in cell-cell recognition and signaling.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.