


Product Name: Recombinant Human Hemopexin (C-6His)
Catalog #: PHH0784



概述 (Summary)

英文全称	Hemopexin
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Hemopexin is produced by our Mammalian expression system and the target gene encoding Thr24-His462 is expressed with a 6His tag at the C-terminus.
Accession #	P02790
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	50.1 KDa
蛋白形态 (Form)	Supplied as a 0.2 μm filtered solution of 20mM MES, 150mM NaCl, pH 5.5.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	

电泳图 (SDS-PAGE image)



背景 (Background)

分子别名 (Alternative Names) Hemopexin;Hpx;Hpxn

Product Name: Recombinant Human Hemopexin (C-6His)
Catalog #: PHH0784



背景介绍 (References)

Hemopexin (HPX) is plasma glycoprotein belongs to the family of the acute-phase proteins whose synthesis is induced after an inflammatory event. Hemopexin with two four-bladed beta -propeller folds has been found in other proteins including collagenases and provides sites for protein-protein interactions. The liver is the major synthesizing organ. Hemopexin participates in maintaining and recycling the iron pool by utilizing its high binding affinity toward heme composed of protoporphyrin IX and iron. It also functions in preventing oxidation caused by heme after hemolysis. Hydrophobic heme molecules can intercalate into lipid membranes and participate in the oxidation of lipid membrane components through the Fenton reaction resulting in lipid peroxidation. Hemopexin undergoes a conformational change upon the binding of heme. The conformational change allows hemopexin to interact with a specific receptor, forming a complex which is then internalized. Heme concentrations in plasma increase after hemolysis, which is associated with several pathological conditions such as reperfusion injury and ischemia.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.