产品名称：Recombinant Human LILRB5 (C-6His)
产品货号：PHH1088

概述（Summary）

英文全称	LILRB5/LIR8/CD85c
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	Recombinant Human Leukocyte Immunoglobulin-like Receptor Subfamily B Member 5 is produced by our Mammalian expression system and the target gene encoding Gly24-His456 is expressed with a 6His tag at the C-terminus.
Accession #	O75023
表达宿主（Host）	Human Cells
种属（Species）	Human
预测分子量（Predicted MW）	47.8 KDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
运输方式（Shipping）	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图（SDS-PAGE image）

产品名称：Recombinant Human LILRB5 (C-6His)
产品货号：PHH1088



背景（Background）

分子别名（Alternative Names）

Leukocyte immunoglobulin-like receptor subfamily B member 5; CD85 antigen-like family member C; Leukocyte immunoglobulin-like receptor 8; LIR-8; CD85c; LILRB5; LIR8

背景介绍（References）

Human Leukocyte Immunoglobulin-like Receptor Subfamily B Member 5 (LILRB5/CD85c/LIR-8) belongs to a family of transmembrane glycoproteins that negatively regulate immune cell activation. Mature human LIR-8 consists of a 435 amino acid (aa) extracellular domain with four Iglike domains, a 21 aa transmembrane segment, and a 111 aa cytoplasmic domain with two immunoreceptor tyrosine-based inhibitory motifs (ITIM). Alternative splicing of human LIR-8 generates an isoform that lacks the second Ig-like domain. LIR-8 is expressed on NK cells and in the tryptic granules of mast cells. Following cell activation and degranulation, it is present on the mast cell surface. Activated mast cells may also release soluble forms of LIR-8.

注意事项（Note）

For research use only .