


概述 (Summary)

英文全称	LILRB5/LIR8/CD85c
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Leukocyte Immunoglobulin-like Receptor Subfamily B Member 5 is produced by our Mammalian expression system and the target gene encoding Gly24-His456 is expressed with a 6His tag at the C-terminus.
Accession #	O75023
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	47.8 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human LILRB5 (C-6His)
Catalog #: PHH1088



背景 (Background)

分子别名 (Alternative Names)

Leukocyte immunoglobulin-like receptor subfamily B member 5; CD85 antigen-like family member C; Leukocyte immunoglobulin-like receptor 8; LIR-8; CD85c; LILRB5; LIR8

背景介绍 (References)

Human Leukocyte Immunoglobulin-like Receptor Subfamily B Member 5 (LILRB5/CD85c/LIR-8) belongs to a family of transmembrane glycoproteins that negatively regulate immune cell activation. Mature human LIR-8 consists of a 435 amino acid (aa) extracellular domain with four Iglike domains, a 21 aa transmembrane segment, and a 111 aa cytoplasmic domain with two immunoreceptor tyrosine-based inhibitory motifs (ITIM). Alternative splicing of human LIR-8 generates an isoform that lacks the second Ig-like domain. LIR-8 is expressed on NK cells and in the tryptic granules of mast cells. Following cell activation and degranulation, it is present on the mast cell surface. Activated mast cells may also release soluble forms of LIR-8.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.