


Product Name: Recombinant Human NGAL (C-6His, Human Cells)
Catalog #: PHH1090

概述 (Summary)

英文全称	NGAL/Lipocalin-2/LCN2/Neutrophil gelatinase-associated lipocalin/p25
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Neutrophil Gelatinase-associated Lipocalin is produced by our Mammalian expression system and the target gene encoding Gln21-Gly198 is expressed with a 6His tag at the C-terminus.
Accession #	P80188
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	21.6 KDa
蛋白形态 (Form)	Supplied as a 0.2 μm filtered solution of PBS, 50% Glycerol, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	

电泳图 (SDS-PAGE image)



背景 (Background)

分子别名 (Alternative Names) Neutrophil gelatinase-associated lipocalin; NGAL; 25 kDa alpha-2-

Product Name: Recombinant Human NGAL (C-6His, Human Cell)
Catalog #: PHH1090



背景介绍 (References)

microglobulin-related subunit of MMP-9; Lipocalin-2; Oncogene 24p3; Siderocalin LCN2; p25; HNL; NGAL

LCN2 is iron-trafficking protein involved in multiple processes such as apoptosis, innate immunity and renal development. LCN2 binds iron through association with 2,5-dihydroxybenzoic acid (2,5-DHBA), a siderophore that shares structural similarities with bacterial enterobactin, and delivers or removes iron from the cell, depending on the context. LCN2 is involved in apoptosis due to interleukin-3 (IL3) deprivation: iron-loaded form increases intracellular iron concentration without promoting apoptosis, while iron-free form decreases intracellular iron levels, inducing expression of the proapoptotic protein BCL2L11/BIM, resulting in apoptosis. LCN2 is involved in innate immunity, possibly by sequestering iron, leading to limit bacterial growth.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.