


概述 (Summary)

英文全称	LRRTM2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Leucine-Rich Repeat Transmembrane Neuronal Protein 2 is produced by our Mammalian expression system and the target gene encoding Cys34-Arg422 is expressed with a 6His tag at the C-terminus.
Accession #	O43300
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	45.55 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human LRRTM2 (C-6His)
Catalog #: PHH1102



背景 (Background)

分子别名 (Alternative Names)

Leucine-Rich Repeat Transmembrane Neuronal Protein 2; Leucine-Rich Repeat Neuronal 2 Protein; LRRTM2; KIAA0416; LRRN2

背景介绍 (References)

Leucine-Rich Repeat Transmembrane Neuronal Protein 2 (LRRTM2) is a single-pass type I membrane protein that belongs to the LRRTM family. It contains ten LRR (leucine-rich) repeats, one LRRCT domain, and one LRRNT domain. LRRTM2 is expressed in neuronal tissues, and it interacts with DLG4 and NRXN1. LRRTM2 has been suggested to be involved in the development and maintenance of excitatory synapses in the vertebrate nervous system. LRRTM2 also regulates the surface expression of AMPA receptors. LRRTM2 acts as a ligand for the presynaptic receptors NRXN1-A and NRXN1-B.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.