


概述 (Summary)

英文全称	Myelin peripheral protein/MPZ
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Myelin Protein P0 is produced by our Mammalian expression system and the target gene encoding Ile30-Arg153 is expressed with a 6His tag at the C-terminus.
Accession #	P25189
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	15.2 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human MPZ (C-6His)
Catalog #: PHH1184



背景 (Background)

分子别名 (Alternative Names)

背景介绍 (References)

Myelin Protein P0; Myelin Peripheral Protein; MPP; Myelin Protein Zero; MPZ

Myelin Protein P0 (MPZ) is a single-pass type I membrane glycoprotein which belongs to the myelin P0 protein family. MPZ contains one Ig-like V-type (immunoglobulin-like) domain, absent in the central nervous system. MPZ is a major component of the myelin sheath in peripheral nerves. It is postulated that MPZ is a structural element in the formation and stabilisation of peripheral nerve myelin, holding its characteristic coil structure together by the interaction of its positively-charged domain with acidic lipids in the cytoplasmic face of the opposed bilayer, and by interaction between hydrophobic globular of adjacent extracellular domains. Defects in MPZ associated with Charcot-Marie-Tooth disease and Dejerine-Sottas disease.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.