


概述 (Summary)

英文全称	Neuroplastin
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Neuroplastin is produced by our Mammalian expression system and the target gene encoding Gln29-His220 is expressed with a 6His tag at the C-terminus.
Accession #	Q9Y639
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	22.69 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human NPTN (C-6His)
Catalog #: PHH1216



背景 (Background)

分子别名 (Alternative Names)

Neuroplastin; Stromal Cell-Derived Receptor 1; SDR-1; NPTN; SDFR1; SDR1

背景介绍 (References)

Neuroplastin (NPTN) is a 52-57 kDa member of the Ig-superfamily. Neuroplastin likely serves as a cell adhesion molecule, and is widely expressed in multiple tissues. Human Neuroplastin is 282 amino acids that contains two Ig-like domains and a 38 aa cytoplasmic region. Probable homophilic and heterophilic cell adhesion molecule involved in long term potentiation at hippocampal excitatory synapses through activation of p38MAPK. Neuroplastin may also regulate neurite outgrowth by activating the FGFR1 signaling pathway. It may play a role in synaptic plasticity.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.