


概述 (Summary)

英文全称	NgR3/NgRH2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Nogo-66 Receptor-related Protein 3/Reticulon-4 Receptor-like 1 is produced by our Mammalian expression system and the target gene encoding Cys25-Ala419 is expressed with a 6His tag at the C-terminus.
Accession #	Q86UN2
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	45.49 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human NgR3 (C-6His)
Catalog #: PHH1226



背景 (Background)

分子别名 (Alternative Names)

Reticulon-4 Receptor-Like 1; Nogo Receptor-Like 2; Nogo-66 Receptor Homolog 2; Nogo-66 Receptor-Related Protein 3; NgR3; RTN4RL1; NGRH2; NGRL2

背景介绍 (References)

Nogo-66 Receptor-Related Protein 3 (NgR3) has primary structures with NgR2 (NgRH1, NgRL3) and biochemical properties that are homologous to Nogo-66 receptor (NgR), and constitute a novel neuronal receptor protein family. NgR is GPI-anchored and contains eight leucine-rich repeats (LRR), it is the neuronal receptor for the myelin-associated proteins Nogo-A, OMgp (oligodendrocyte myelin glycoprotein), and MAG (myelin-associated glycoprotein) and mediates the inhibition of CNS axonal regeneration both in vitro and in vivo. NgR2 and NgR3 have similar structure and distinct but overlapping expression versus NgR. NgR2 can be metalloproteinase-cleaved to release a soluble ectodomain. NgR2 has also been shown to bind MAG, but ligands for NgR3 have not yet been determined. Mature human NgR3 shares 88%, 88%, 48% and 44% amino acid identity with mature mouse NgR3, rat NgR3, human NgRH1 and NgR, respectively.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.