Product Name: Recombinant Human PCDH10 (C-Fc)

Catalog #: PHH1271



概述 (Summary)

英文全称 PCDH10/protocadherin-10

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/μg as determined by LAL test.

蛋白构建 (Construction) Recombinant Human Protocadherin-10 is produced by our Mammalian

expression system and the target gene encoding Gln19-Thr715 is

expressed with a human IgG1 Fc tag at the C-terminus.

Accession # Q9P2E7

蛋白标签 (Tag)

表达宿主 (Host) Human Cells

种属 (Species)Human预测分子量 (Predicted MW)102.4 KDa

蛋白形态 (Form) Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

稳定性&储存 (Stability &Storage) Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to

is not recommended to reconstitute to a concentration less than 100µg/ml.

reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize

freeze-thaw cycles.


电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human PCDH10 (C-Fc)

Catalog #: PHH1271





背景 (Background)

分子別名 (Alternative Names) 背景介绍 (References) Protocadherin-10; PCDH10; KIAA1400

Protocadherin-10 (PCDH10) is a single-pass type I membrane protein that contains six extracellular cadherin domains, one transmembrane domain and one cytoplasmic tail differing from those of the classical cadherins. As a potential calcium-dependent cell-adhesion neuronal receptor, it may plays a role in the establishment and function of specific cell-cell connections in the brain. PCDH10 moderately expressed in all regions of the brain examined, as well as in testis and ovary, and low expression in all other tested tissues.

注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.