


概述 (Summary)

英文全称	PDGF R beta/PDGFRB/PDGFR β /CD140b
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/ μ g as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Platelet-Derived Growth Factor Receptor Beta is produced by our Mammalian expression system and the target gene encoding Leu33-Phe530 is expressed with a 6His tag at the C-terminus.
Accession #	AAH32224.1
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	57.17 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μ m filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at $\leq -70^{\circ}\text{C}$, stable for 6 months after receipt. Store at $\leq -70^{\circ}\text{C}$, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human PDGFRB (C-6His)
Catalog #: PHH1289



背景 (Background)

分子别名 (Alternative Names)

Platelet-Derived Growth Factor Receptor Beta; PDGF-R-Beta; PDGFR-Beta; Beta Platelet-Derived Growth Factor Receptor; Beta-Type Platelet-Derived Growth Factor Receptor; CD140 Antigen-Like Family Member B; Platelet-Derived Growth Factor Receptor 1; PDGFR-1; CD140b; PDGFRB; PDGFR; PDGFR1

背景介绍 (References)

Platelet-Derived Growth Factor Receptor β (PDGFR- β) is a member of the protein kinase superfamily and CSF-1/PDGF receptor subfamily. The PDGF family consists of PDGF-A, -B, -C and -D, which form either homo- or heterodimers (PDGF-AA, -AB, -BB, -CC, -DD). The four PDGFs are inactive in their monomeric forms. The PDGFs bind to the protein tyrosine kinase receptors PDGF receptor- α and - β . These two receptor isoforms dimerize upon binding the PDGF dimer, leading to three possible receptor combinations, namely - $\alpha\alpha$, - $\beta\beta$ and - $\alpha\beta$. The extracellular region of the PDGF receptor- β consists of five immunoglobulin-like domains while the intracellular part is a tyrosine kinase domain. In addition to being a potent mitogen for cells of mesenchymal origin, PDGF has also been shown to be a potent chemoattractant for mesenchymal cells, mononuclear cells, and neutrophils and has been reported to be important in the modification of cellular matrix constituents.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.