


概述 (Summary)

英文全称	Periostin(POSTN)
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Osteoblast Specific Factor 2 is produced by our Mammalian expression system and the target gene encoding Asn22-Gln779 is expressed with a 6His tag at the C-terminus.
Accession #	Q15063-2
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	85.6 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human Periostin (C-6His)
Catalog #: PHH1307



背景 (Background)

分子别名 (Alternative Names)

Osteoblast-specific factor 2; OSF-2; POSTN; Periostin

背景介绍 (References)

Osteoblast-Specific Factor OSF-2 (POSTN, Periostin, OSF-2) is a secreted, homodimeric protein that belongs to the periostin family of the FAS1 superfamily of molecules. Periostin is a disulfide linked 90kDa bone adhesion protein secreted by osteoblasts and osteoblast-like cell lines and the protein is an attachment agent for osteoblasts. It is a TGF-beta inducible molecule that serves as both an adhesion molecule and tumor suppressor. It is synthesized by smooth muscle cells, fibroblasts and osteoblasts, as well as in the periosteum and periodontal ligament. Periostin functions as a ligand for alpha-V/beta-3 and alpha-V/beta-5 integrins to support adhesion and migration of epithelial cells.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.