# Product Name: Recombinant Human PRKAR1A (C-6His) Enkilife Catalog #: PHH1367

## 概述 (Summary)

**英文全称** PRKAR1A/TSE1

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/μg as determined by LAL test.

蛋白构建 (Construction) Recombinant Human cAMP-dependent Protein Kinase Regulatory Type I-

alpha is produced by our Mammalian expression system and the target gene encoding Glu2-Val381 is expressed with a 6His tag at the C-

terminus.

Accession # P10644

蛋白标签 (Tag)

表达宿主 (Host) Human Cells 种属 (Species) Human 预测分子量 (Predicted MW) 44 KDa

蛋白形态 (Form) Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

稳定性&储存 (Stability &Storage) Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to

is not recommended to reconstitute to a concentration less than 100µg/ml.

reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized


protein in distilled water. Please aliquot the reconstituted solution to minimize

freeze-thaw cycles.

# 电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





# 背景 (Background)

分子別名 (Alternative Names) 背景介绍 (References) Tissue-specific extinguisher 1;TSE1

cAMP-dependent protein kinase type I-alpha regulatory subunit is an enzyme that in humans is encoded by the PRKAR1A gene. cAMP is a signaling molecule important for a variety of cellular functions. cAMP exerts its effects by activating the cAMP-dependent protein kinase A (PKA), which transduces the signal through phosphorylation of different target proteins. Four different regulatory subunits and three catalytic subunits of PKA have been identified in humans. The protein encoded by this gene is one of the regulatory subunits. This protein was found to be a tissue-specific extinguisher that down-regulates the expression of seven liver genes in hepatoma x fibroblast hybrids.

## 注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.