


概述 (Summary)

英文全称	RANK/TNFRSF11A/CD265
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Receptor Activator of NF-kappa-B is produced by our Mammalian expression system and the target gene encoding Ile30-Pro212 is expressed with a 6His tag at the C-terminus.
Accession #	Q9Y6Q6
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	21.1 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human RANK (C-6His)
Catalog #: PHH1411



背景 (Background)

分子别名 (Alternative Names)

CD265; ODFR; TNFRSF11A; TRANCE R; CD265; CD265 antigen; FEO; ODFROSTS; OFE; OPTB7; PDB2; RANK1; Receptor activator of NF- κ B; receptor activator of nuclear factor-kappa B; TRANCER; tumor necrosis factor receptor superfamily member 11A

背景介绍 (References)

Receptor Activator of Nuclear Factor κ B (RANK), also known as CD265, TRANCE Receptor or TNFRSF11A, is member of the tumor necrosis factor receptor (TNFR) molecular superfamily. RANK is the receptor for RANK-Ligand (RANKL) and part of the RANK/RANKL/OPG signaling pathway that regulates osteoclast differentiation and activation. It plays a vital role in bone remodeling and repair, immune cell function, lymph node development, thermal regulation, and mammary gland development. RANK is constitutively expressed in skeletal muscle, thymus, liver, colon, small intestine, adrenal gland, osteoclast, mammary gland epithelial cells, prostate, vascular cell, and pancreas.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.