


产品名称: Recombinant Human RSPO1 (C-6His)
产品货号: PHH1448



概述 (Summary)

| | |
|-----------------------------|---|
| 英文全称 | R-spondin-1/RSPO1/Rspondin-1 |
| 纯度 (Purity) | Greater than 95% as determined by reducing SDS-PAGE |
| 内毒素 (Endotoxin level) | <0.01 EU/μg as determined by LAL test. |
| 蛋白构建 (Construction) | Recombinant Human R-spondin-1 is produced by our Mammalian expression system and the target gene encoding Ser21-Ala263 is expressed with a 6His tag at the C-terminus. |
| Accession # | Q2MKA7 |
| 表达宿主 (Host) | Human Cells |
| 种属 (Species) | Human |
| 预测分子量 (Predicted MW) | 27.8 KDa |
| 制剂 (Form) | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4. |
| 运输方式 (Shipping) | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. |
| 稳定性&储存 (Stability &Storage) | Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months. |
| 复溶 (Reconstitution) | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

电泳图 (SDS-PAGE image)



产品名称: Recombinant Human RSPO1 (C-6His)
产品货号: PHH1448



背景 (Background)

分子别名 (Alternative Names)

RSPO1; R-spondin1; RP11-566C13.1; CRISTIN3; FLJ40906; RSPO Rspo1; R-spondin; Rspondin; RP23-325M14.2; Roof plate-specific spondin-1

背景介绍 (References)

RSPO1 is a secreted protein, containing 2 FU(furin-like) repeats and 1 TSP type-1 domain and belonging to the R-spondin family. RSPO1 is required for the early development of gonads, regardless of sex. It has been found in mice only eleven days after fertilization. To induce cell proliferation, it acts synergistically with WNT4. They help stabilize β catenin, which activates downstream targets. RSPO1 is necessary in female sex development. It augments the WNT/ β catenin pathway to oppose male sex development. In critical gonadal stages, between six and nine weeks after fertilization, the ovaries upregulate it while the testes downregulate it. RSPO1 can potentially aid in the treatment of mucositis, which is characterized by inflammation of the oral cavity. This unfortunate condition often accompanies chemotherapy and radiation in cancer patients with head and neck tumors.

注意事项 (Note)

For research use only .