


概述 (Summary)

英文全称	sCD4/T-cell surface glycoprotein CD4
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human T-cell Surface Glycoprotein CD4 is produced by our Mammalian expression system and the target gene encoding Lys26-Trp390 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	P01730
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	67.6 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, 1mM EDTA, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human sCD4 (C-Fc)
Catalog #: PHH1466



背景 (Background)

分子别名 (Alternative Names)

背景介绍 (References)

T-cell surface glycoprotein CD4; T-cell surface antigen T4/Leu-3; CD4; Scd4

CD4 is an approximately 55 kDa type I transmembrane glycoprotein that is expressed predominantly on thymocytes and a subset of mature T lymphocytes. It is a standard phenotype marker for the identification of T cell populations. Mature human CD4 consists of a 371 amino acid extracellular region containing four immunoglobulin-like domains, a 22 aa transmembrane segment, and a 40 aa cytoplasmic domain. CD4 is expressed along with CD8 on double positive T cells during their development in the thymus. CD4 binds directly to MHC class II molecules on antigen presenting cells (10). This interaction contributes to the formation of the immunological synapse which is focused around the TCR-MHC class II-antigenic peptide interaction. CD4 also functions as a chemotactic receptor for IL-16 and, in human, as a co-receptor for the gp120 surface glycoprotein of HIV-1.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.