

Product Name: Recombinant Human SHMT1 (C-6His)
Catalog #: PHH1492




概述 (Summary)

英文全称	Serine hydroxymethyltransferase/SHMT1
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Serine Hydroxymethyltransferase Cytosolic is produced by our Mammalian expression system and the target gene encoding Met3-Phe483 is expressed with a 6His tag at the C-terminus.
Accession #	AAH07979.1
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	53.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, 1mM EDTA, 5% Trehalose, 5% Mannitol, 0.02% Tween80, pH 6.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human SHMT1 (C-6His)
Catalog #: PHH1492



背景 (Background)

分子别名 (Alternative Names)

Serine Hydroxymethyltransferase Cytosolic; SHMT; Glycine Hydroxymethyltransferase; Serine Methylase; SHMT1

背景介绍 (References)

Serine Hydroxymethyltransferase Cytosolic (SHMT1) is a member of the SHMT family. SHMT1 is a cytoplasmic protein and exists as a homotetramer. SHMT1 catalyzes the reversible conversion of serine and tetrahydrofolate to glycine and 5,10-methylene tetrahydrofolate. This reaction provides one carbon unit for the synthesis of methionine, thymidylate, and purines in the cytoplasm. A reduction in SHMT1 levels would result in less glycine that could affect the nervous system by acting as an agonist to the NMDA receptor and this could be a mechanism behind Smith-Magenis syndrome.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.