

Product Name: Recombinant Human TXNDC15 (C-6His)
Catalog #: PHH1751




概述 (Summary)

英文全称	TXNDC15/Thioredoxin domain-containing protein 15
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Thioredoxin Domain-Containing Protein 15 is produced by our Mammalian expression system and the target gene encoding Val33-Ser321 is expressed with a 6His tag at the C-terminus.
Accession #	Q96J42
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	32.5 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human TXNDC15 (C-6His)
Catalog #: PHH1751



背景 (Background)

分子别名 (Alternative Names)

Thioredoxin domain-containing protein 15;C5orf14;UNQ335/PRO534

背景介绍 (References)

Thioredoxin domain-containing protein 15(TXNDC15) is a single-pass type I membrane protein. Mature Human TXNDC15 consists of a 289 amino acid (aa) extracellular region (ECD) with one thioredoxin domain, a 21 aa transmembrane domain, and a 18 aa cytoplasmic region. It has 2 isoforms produced by alternative splicing. Thioredoxins comprise a family of small proteins that, by catalyzing the oxidation of disulfide bonds, participate in redox reactions throughout the cell. Proteins that contain thioredoxin domains do not necessarily convey the oxidative properties of thioredoxins, but generally function as disulfide isomerases that enzymatically rearrange disulfide bonds found in various proteins.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.