


概述 (Summary)

英文全称	V-set and transmembrane domain-containing protein 1/VSTM1
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human V-set And Transmembrane Domain-Containing Protein 1 is produced by our Mammalian expression system and the target gene encoding Tyr17-Thr135 is expressed with a 6His tag at the C-terminus.
Accession #	Q6UX27
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	14.6 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human VSTM1 (C-6His)
Catalog #: PHH1823



背景 (Background)

分子别名 (Alternative Names)

V-set and transmembrane domain-containing protein 1;SIRL-1;Signal inhibitory receptor on leukocytes-1;VSTM1

背景介绍 (References)

V-set and transmembrane domain-containing protein 1 is a single-pass membrane protein. VSTM1 Contains 1 Ig-like V-type domain, in humans is encoded by the VSTM1 gene. It expressed on myeloid (neutrophils, eosinophils and monocytes) but not on lymphoid cells. It behaves as a cytokine, promoting IL17A secretion by CD4+ T-cells, and differentiation and activation of IL17 producing helper T-cells (TH17). Inhibitory immune receptor involved in the regulation of phagocytes.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.