

Product Name: Recombinant Human Follistatin 288 (C-6His)
Catalog #: PHH1897




概述 (Summary)

英文全称	Follistatin/Follistatin 288/FST
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Follistatin/FST is produced by our Mammalian expression system and the target gene encoding Gly30-Asn317 is expressed with a 6His tag at the C-terminus.
Accession #	P19883
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	32.4 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human Follistatin 288 (C-6His)
Catalog #: PHH1897



背景 (Background)

分子别名 (Alternative Names)

follistatin isoform FST317; Follistatin; FS; FSActivin-binding protein; FST

背景介绍 (References)

Follistatin 288 is a secreted glycoprotein that was first identified as a follicle-stimulating hormone inhibiting substance in ovarian follicular fluid. Human follistatin 288 cDNA encodes a 317 amino acid (aa) protein with a 29/aa signal sequence, and a 288 aa mature region. Follistatin shows the highest affinity for activins due to its extended configuration. Genetic deletion of follistatin in mice, or expression of only the Follistatin form, is perinatally lethal due to defects of lung, skin and musculoskeletal system. Follistatins also regulate hematopoietic stem cell adhesion to fibronectin via FS2.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.