


概述 (Summary)

英文全称	MMP-8/Matrix metalloproteinase-8
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Neutrophil Collagenase is produced by our Mammalian expression system and the target gene encoding Phe21-Gly467 is expressed with a 10His tag at the C-terminus. The proenzyme needs to be activated by APMA for an activated form.
Accession #	P22894
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	52.8 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human MMP-8 (C-10His)
Catalog #: PHH1905



背景 (Background)

分子别名 (Alternative Names)

Neutrophil collagenase; Matrix metalloproteinase-8; MMP-8; PMNL collagenase; PMNL-CL; MMP8; CLG1

背景介绍 (References)

Matrix metalloproteinases (MMPs) are a family of zinc and calcium dependent endopeptidases with the combined ability to degrade all the components of the extracellular matrix. MMP8(neutrophil collagenase) is expressed in neutrophils, where it is stored in specific granules. MMP8 release from the neutrophils is stimulated by various factors such as interleukins 1 and 8, TNF- α and GM-CSF. MMP8 is capable of cleaving types I, II and III triple-helical collagen, gelatin peptides, fibronectin, proteoglycans, aggrecan, serpins, β -casein and peptides such as angiotensin and substance P. In addition to its function in phagocytosis, MMP8 has a high capacity for infiltrating connective tissue, and is implicated in the breakdown of the extracellular matrix in diseases such as rheumatoid arthritis. MMP8 is heavily glycosylated.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.