


概述 (Summary)

英文全称	CREG/CREG1/Protein CREG1
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Cellular Repressor Of E1A-stimulated Genes 1 is produced by our Mammalian expression system and the target gene encoding Arg32-Gln220 is expressed with a 6His tag at the C-terminus.
Accession #	O75629
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	21.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human CREG (C-6His)
Catalog #: PHH1924



背景 (Background)

分子别名 (Alternative Names)

CREG1; cellular repressor of E1A-stimulated genes; cellular repressor of E1A-stimulated genes 1CREG; protein CREG1

背景介绍 (References)

Cellular repressor of E1A genes (CREG) is an evolutionarily conserved lysosomal protein, and an important new factor in regulating tissue homeostasis that has been shown to antagonize injury of tissues or cells. CREG contains three mannose 6-phosphate (M6P) markers, and depends on interactions with M6P receptors for efficient delivery to lysosomes, which is implicated in the regulation of lysosomal functions. This protein shares limited sequence similarity with E1A and binds both the general transcription factor TBP and the tumor suppressor pRb in vitro. CREG plays an important role in the control of cell growth and differentiation. It has been shown that CREG antagonizes transcriptional and cellular transformation by the adenoviral E1A oncoprotein, induces differentiation while attenuating cellular proliferation, regulates the levels of the signaling kinases ERK1/2, and mediates glucocorticoid-induced proliferation of ileal epithelial cells. CREG is widely expressed in adult tissues, such as the brain, heart, lungs, liver, intestines and kidneys in mice, but is not markedly expressed in pluripotent embryonic stem cells.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.