

产品名称: Recombinant Human OX40L (C-mFc)
产品货号: PHH2044




概述 (Summary)

英文全称	OX40 Ligand/OX40L/TNFSF4/CD252/Tumor necrosis factor ligand superfamily member 4
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human OX40 Ligand is produced by our Mammalian expression system and the target gene encoding Gln51-Leu183 is expressed with a mouse IgG1 Fc tag at the C-terminus.
Accession #	P23510
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	42.2 KDa
制剂 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

产品名称: Recombinant Human OX40L (C-mFc)
产品货号: PHH2044



背景 (Background)

分子别名 (Alternative Names)

Tumor necrosis factor ligand superfamily member 4; Glycoprotein Gp34; OX40 ligand; OX40L; TAX transcriptionally-activated glycoprotein 1; TNFSF4; CD252; TXGP1

背景介绍 (References)

Tumor necrosis factor ligand superfamily member 4(TNFSF4/OX40L) is a single-pass type II membrane protein.OX40L is expressed on the surface of activated B cells, T cells, dendritic cells and endothelial cells. OX40L binds to OX40 (CD134), a member of the TNF receptor superfamily that is expressed predominantly on activated CD4+ T cells. OX40-OX40L co-stimulates signal to promote the survival and proliferation of activated CD4+ T cells and prolong the immune response. It involved in T-cell proliferation and cytokine production. Additional, it has been found association with systemic lupus erythematosus, no association with occurrence of atherosclerosis.

注意事项 (Note)

For research use only .