


概述 (Summary)

英文全称	CD155/PVR/Poliovirus Receptor/Nectin-Like Protein 5/NECL-5/PVS
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Poliovirus Receptor is produced by our Mammalian expression system and the target gene encoding Trp21-Asn343 is expressed with a mouse IgG1 Fc tag at the C-terminus.
Accession #	P15151
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	61.7 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)



Product Name: Recombinant Human PVR (C-mFc)
Catalog #: PHH2061



背景 (Background)

分子别名 (Alternative Names)

Poliovirus Receptor; Nectin-Like Protein 5; NECL-5; CD155; PVR; PVS

背景介绍 (References)

Poliovirus Receptor (PVR) is a 70 kDa type I transmembrane single-span glycoprotein that belongs to the nectin-like (Nect) family and was originally identified based on its ability to mediate the cell attachment and entry of poliovirus (PV), an etiologic agent of the central nervous system disease poliomyelitis. PVR contains three Ig-like extracellular domains, a transmembrane segment, and a cytoplasmic tail. The normal cellular function of PVR maybe the involvement of intercellular adhesion between epithelial cells. Alternate splicing of the PVR mRNA yields four different isoforms (α , β , γ , and δ) with identical extracellular domains.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.