


概述 (Summary)

英文全称	LILRB2/ILT4/CD85d/Lir-2/LIR2/MIR10/Leukocyte Ig-Like Receptor B2/Leukocyte Immunoglobulin-Like Receptor Subfamily B Member 2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Leukocyte Immunoglobulin-Like Receptor Subfamily B Member 2 is produced by our Mammalian expression system and the target gene encoding Gln22-His458 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	AAH36827.1
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	74.5 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human LILRB2 (C-Fc)
Catalog #: PHH2145



背景 (Background)

分子别名 (Alternative Names)

Leukocyte Immunoglobulin-Like Receptor Subfamily B Member 2; LIR-2; Leukocyte Immunoglobulin-Like Receptor 2; CD85 Antigen-Like Family Member D; Immunoglobulin-Like Transcript 4; ILT-4; Monocyte/Macrophage Immunoglobulin-Like Receptor 10; MIR-10; CD85d; LILRB2; ILT4; LIR2; MIR10

背景介绍 (References)

Members of the immunoglobulin-like transcript (ILT) family are activating and inhibitory immunoreceptors whose genes are located same locus that encodes killer cell Ig-like receptors (KIR). Leukocyte Immunoglobulin-Like Receptor Subfamily B Member 2 (LIR-2) is a type I transmembrane protein. LIR-2 is expressed primarily on monocytes and dendritic cells (DC). Human LIR-2 is produced as a 598 amino acid precursor including a 21 aa signal sequence, a 440 aa extracellular domain (ECD), a 21 aa transmembrane segment, and a 116 aa cytoplasmic domain. LIR-2 binds to Classical MHC I proteins. Ligation of LIR-2 includes Tyr phosphorylation within its cytoplasmic ITIMs, a requirement for association with SHP-1. LIR-2 mediates tolerogenic DC-induced CD4⁺ T cell energy in vitro and in vivo.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.