


概述 (Summary)

英文全称	CD82
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/ μ g as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human CD82 Antigen is produced by our Mammalian expression system and the target gene encoding Gly103-Gln225 is expressed with a human IgG1 Fc tag at the N-terminus.
Accession #	P27701
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	40.3 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at $\leq -70^{\circ}\text{C}$, stable for 6 months after receipt. Store at $\leq -70^{\circ}\text{C}$, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human CD82 (N-Fc)
Catalog #: PHH2153



背景 (Background)

分子别名 (Alternative Names)

CD82 antigenC33; CD82 molecule; CD82; IA4; IA4C33 antigen; Inducible membrane protein R2; KAI1; KAI1GR15; ST6; ST6tspan-27; Suppressor of tumorigenicity 6 protein; Tetraspanin-27; TSPAN27; Tspan-27; TSPAN274F9

背景介绍 (References)

CD82 antigen, also known as Kai-1, is a widely expressed palmitoylated molecule of the tetraspanin superfamily. KAI1/CD82 is localized on cell membrane and form interactions with other tetraspanins, integrins and chemokines which are respectively responsible for cell migration, adhesion and signaling. CD82/Kai-1 is a component of the promiscuous TIMP-1 interacting protein complex on the cell surface of human adenocarcinoma cells and gives insight into tumorigenic metastatic potential. CD82/Kai-1 suppresses EMT in prostate cancer cells adhered to fibronectin leading to reduced cell migration and invasiveness. CD82/Kai-1 function is important for muscle stem cell function in muscular disorders. Overexpression of CD82/Kai-1 suppresses growth, migration and invasion of oral cancer cells and may be considered as a potential therapeutic target in oral cancer.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.