

产品名称: Recombinant Human Otolin-1 (C-6His)
产品货号: PHH2211




概述 (Summary)

英文全称	Otolin-1
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/ μ g as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Otolin-1 is produced by our Mammalian expression system and the target gene encoding Lys24-Pro477 is expressed with a 6His tag at the C-terminus.
Accession #	A6NHN0
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	47.7 KDa
制剂 (Form)	Lyophilized from a 0.2 μ m filtered solution of PBS, 5% Trehalose, pH 7.4.
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at $\leq -20^{\circ}\text{C}$, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8 $^{\circ}\text{C}$ for 2-7 days. Aliquots of reconstituted samples are stable at $\leq -20^{\circ}\text{C}$ for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

产品名称: Recombinant Human Otolin-1 (C-6His)
产品货号: PHH2211



背景 (Background)

分子别名 (Alternative Names)

OTOL1; otolin 1; Otolin-1; C1qTNF15

背景介绍 (References)

Otolin (OTOL1), also known as C1qTNF15, is an approximately 65 kDa protein found in the otoconial membrane lining the cochlea and vestibular labyrinth of the inner ear. Collagen-like protein specifically expressed in the inner ear, which provides an organic scaffold for otoconia, a calcium carbonate structure in the saccule and utricle of the ear. It associates into multimers and disulfide-linked oligomers and also associates with other otoconial proteins including and Otoconin-90 (also known as PLA2G2A, PLA2L, and phospholipase A2 homolog) and Cerebellin-1. It is extensively glycosylated and has multiple hydroxylated proline residues in the collagenous regions.

注意事项 (Note)

For research use only .