


概述 (Summary)

英文全称	ILDR2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Immunoglobulin-like Domain-containing Receptor 2 is produced by our Mammalian expression system and the target gene encoding Leu21-Glu186 is expressed with a 6His tag at the C-terminus.
Accession #	Q71H61
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	19.5 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human ILDR2 (C-6His)
Catalog #: PHH2325



背景 (Background)

分子别名 (Alternative Names)

Angulin-3; C1orf32; Dbm1; DJ782G3.1; ILDR2; immunoglobulin-like domain containing receptor 2; LISCH-Like

背景介绍 (References)

ILDR2 is a member of the B7-like family of proteins that regulate T cell activity, is also a known endoplasmic reticulum molecule that regulates lipid homeostasis. The human ILDR2 luminal domain shares a 99% and 98% homology with the mouse and rat respectively. The human gene encoding ILDR2 is located in a region on Chr1q23–25 that has been associated with type 2 diabetes. ILDR2 plays critical roles in hepatic clearance of lipoproteins and in lipid homeostasis. ILDR2 regulates human dendritic cells (DC2 cells, a subpopulation of polarized DCs that promotes Th2 differentiation). Recent publications reported that ILDR2 displayed negative regulatory functions on human and mouse T cells in various experimental systems. Fusion protein of ILDR2 luminal domain with an Fc fragment, displays therapeutic effects in collagen-induced arthritis (CIA), a mouse model of rheumatoid arthritis (RA). ILDR2 represents a novel B7-like ligand that exerts negative immune modulation via interaction with a putative counterpart receptor expressed on activated T cells.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.