Product Name: Recombinant Human IL-17RB (C-Fc)

Catalog #: PHH2357



概述 (Summary)

英文全称 IL-17RB

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/μg as determined by LAL test.

蛋白构建 (Construction) Recombinant Human Interleukin-17 Receptor B is produced by our

Mammalian expression system and the target gene encoding Arg18-

Gly289 is expressed with a human IgG1 Fc tag at the C-terminus.

Accession # Q9NRM6

蛋白标签 (Tag)

表达宿主 (Host) Human Cells 种属 (Species) Human

预测分子量 (Predicted MW) 56.8 KDa

蛋白形态 (Form) Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

稳定性&储存 (Stability &Storage) Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized

protein in distilled water. Please aliquot the reconstituted solution to minimize

freeze-thaw cycles.


电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human IL-17RB (C-Fc)

Catalog #: PHH2357





背景 (Background)

分子别名 (Alternative Names)

背景介绍 (References)

Evi27; EVI27IL-17B receptor; IL-17 RB; IL-17 receptor B; IL-17 RH1; IL-17ER; IL17Rh1; IL-17Rh1

Interleukin-17B Receptor (IL-17B R), also known as IL-17 Rh1, IL-17E R and EVI27, represents the second receptor of the IL-17 family cytokines to be recognized. IL17RB is the receptor for IL17E, the only member of the IL17 family promoting Th2 reactions. IL17RB is induced on human macrophages by IL4 and enhanced by TGFbeta. Overexpression of IL17RB is associated with poor prognosis and the short survival of breast cancer patients.IL17RB/IL17B signaling triggers a substantial increase in cell growth, proliferation, and migration through the activation of NF-kappaB as well as the up-regulation of the Bcl-2. IL17RB may be the only gene expressed in CD4+ T cells whose transcript measurement is correlated with the variation in IgE level in asthmatics. Diseases associated with IL17RB include Chronic Mucocutaneous Candidiasis and Seborrheic Infantile Dermatitis.

注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.