


概述 (Summary)

英文全称	IL-18RAcP/IL-18RAP/IL-18 Receptor Accessory Protein/IL-1R7
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Interleukin-18 receptor accessory protein is produced by our Mammalian expression system and the target gene encoding Phe20-Arg356 is expressed with a human IgG1 Fc, 6His tag at the C-terminus.
Accession #	O95256
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	66.4 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human IL-18RAcP (C-Fc-6His)
Catalog #: PHH2365



背景 (Background)

分子别名 (Alternative Names)

Interleukin-18 receptor accessory protein; IL18RAP; Accessory protein-like; CD218 antigen-like family member B; CDw218b; IL-1R accessory protein-like; Interleukin-1 receptor 7; Interleukin-18 receptor accessory protein-like; Interleukin-18 receptor beta; CD218b; IL1R7

背景介绍 (References)

IL-18RAP is a single-pass type I membrane protein and contains two Ig-like C2-type domains and one TIR domain, IL18RAP can be induced by IFN- α and IL12 in nature killer cells and T-cells. The coexpression of IL18R1 and IL18RAP is required for the activation of NF-kappa B and JNK in response to IL-18. Defects in IL18RAP are associated with Coeliac disease.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.