


Product Name: Recombinant Human Latent TGF Beta-1&GARP Complex Protein (N-6His, C-3*Strep)
Catalog #: PHH2379

概述 (Summary)

英文全称	Latent TGF Beta-1&GARP Complex Protein
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Latent TGF Beta-1&GARP Complex Protein is produced by our Mammalian expressionsystem and the target gene encoding Leu30-Ser390&His20-Asn627 is expressed with a 6His tag at the N-terminusand, a 3*Strep tag at the C-terminus.
Accession #	P01137&Q14392
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	42.1&71.0 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, 5%Trehalose, 5% Mannitol, 0.05% Tween 80, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human Latent TGF Beta-1&GARP Complex Protein (N-6His, C-3*Strep)
Catalog #: PHH2379



背景 (Background)

分子别名 (Alternative Names)

TGF Beta-1&GARP

背景介绍 (References)

Transforming Growth Factor β -1 (TGF β -1) is a secreted protein which belongs to the TGF- β family. TGF β -1 is abundantly expressed in bone, articular cartilage and chondrocytes and is increased in osteoarthritis (OA). TGF β -1 performs many cellular functions, including the control of cell growth, cell proliferation, cell differentiation and apoptosis. The precursor is cleaved into a latency-associated peptide (LAP) and a mature TGF β -1 peptide. Leucine Rich Repeat Containing 32 (LRRC32), also known as Glycoprotein A Repetitions Predominant (GARP), has been postulated as a novel surface marker of activated T(regs). LRRC32 binds directly to the TGF-beta latency associated peptide (LAP) and tethers latent TGF-beta on the surface of activated Treg cells. The presentation of TGF-beta on Tregs contributes to their ability to suppress naïve T cell proliferation.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.