


Product Name: Recombinant Human CD20 (C-6His-Flag)
Catalog #: PHH2441



概述 (Summary)

英文全称	CD20/B-lymphocyte antigen CD20
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human B-lymphocyte antigen CD20 is produced by our Mammalian expression system and the target gene encoding Met1-Pro297 is expressed with a 6His, Flag tag at the C-terminus. The product is not recommended for cell based experiments.*The product is not recommended for cell based experiments.
Accession #	P11836
蛋白标签 (Tag)	
表达宿主 (Host)	Human cells
种属 (Species)	Human
预测分子量 (Predicted MW)	35.1 kDa
蛋白形态 (Form)	Supplied as a 0.2 μm filtered solution of 50mM HEPES-Na, 150mM NaCl, 0.02%DDM, 0.004%CHS, 5%Glycerol, pH7.5.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	

电泳图 (SDS-PAGE image)



Product Name: Recombinant Human CD20 (C-6His-Flag)
Catalog #: PHH2441



背景 (Background)

分子别名 (Alternative Names)

B-lymphocyte antigen CD20; B-lymphocyte surface antigen B1; Leukocyte surface antigen Leu-16; Membrane-spanning 4-domains subfamily A member 1; Bp35; CD20; MS4A1

背景介绍 (References)

CD20 is a 33-37 kDa non-glycosylated protein expressed on the surface of normal and malignant B lymphocytes, and belongs to the MS4A (membrane-spanning 4-domain family A) protein family. CD20 protein consists of four hydrophobic transmembrane domains, one intracellular and two extracellular domains (large and small loops) with both N- and C- termini residing within the cytosol. CD20 is also known to be physically coupled to major histocompatibility complex class II (MHCII), CD40 molecule, BCR, and the C-terminal src kinase-binding protein (CBP) that interacts with Src kinases such as LYN, FYN, and LCK. CD20 deficiency resulted in a reduced number of circulating memory B cells, reduced isotype switching of Ig, and decreased IgG antibody levels. In agreement with this observation, challenging the patient's primary B cells in vitro using T-dependent and T-independent antigens led to the normal proliferation and secretion of IgM but reduced production of IgG.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.