


概述 (Summary)

英文全称	CD161
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Killer cell lectin-like receptor subfamily B member 1 is produced by our Mammalian expression system and the target gene encoding Gln67-Ser225 is expressed with a 6His tag at the C-terminus.
Accession #	AAI14517.1
蛋白标签 (Tag)	
表达宿主 (Host)	Human cells
种属 (Species)	Human
预测分子量 (Predicted MW)	20.1 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human CD161 (C-6His)
Catalog #: PHH2455



背景 (Background)

分子别名 (Alternative Names)

Killer cell lectin-like receptor subfamily B member 1; KLRB1; CLEC5B; NKR-P1A; CD161; Natural killer cell surface protein P1A; NKR-P1A; HNKRP-1a; C-type lectin domain family 5 member B; KLRB1

背景介绍 (References)

Killer cell lectin-like receptor subfamily B, member 1(KLRB1) is a single-pass type II membrane protein which contains 1 C-type lectin domain. KLRB1 plays an inhibitory role on natural killer (NK) cells cytotoxicity. Activation results in specific acid sphingomyelinase/SMPD1 stimulation with subsequent marked elevation of intracellular ceramide. Activation also leads to AKT1/PKB and RPS6KA1/RSK1 kinases stimulation as well as markedly enhanced T-cell proliferation induced by anti-CD3. It acts as a lectin that binds to the terminal carbohydrate Gal-alpha(1,3)Gal epitope as well as to the N-acetyllactosamine epitope. Binds also to CLEC2D/LLT1 as a ligand and inhibits NK cell-mediated cytotoxicity as well as interferon-gamma secretion in target cells.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.