


概述 (Summary)

英文全称	ALK-1/Activin Receptor-like Kinase 1/ACVRL1
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Activin Receptor-like Kinase 1 is produced by our Mammalian expression system and the target gene encoding Asp23-Pro119 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q61288
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	38.1 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse ALK-1 (C-Fc)
Catalog #: PHM0015



背景 (Background)

分子别名 (Alternative Names)

Serine/threonine-protein kinase receptor R3; SKR3; Activin receptor-like kinase 1; ALK-1; TGF- β superfamily receptor type I; TSR-I; ACVRL1; activin A receptor type II-like 1; activin A receptor

背景介绍 (References)

Activin Receptor-Like Kinase 1 (ALK-1) is a type I cell-surface receptor for the TGF- β superfamily of ligands, which mediates signaling of BMP9 (bone morphogenetic protein) and BMP10. ALK1 signaling is necessary for angiogenesis during embryogenesis, wound healing, and tumor growth. ALK-1 has a high degree of similarity in serine-threonine kinase subdomains, a glycine and serine rich region preceding the kinase-domain, and a C-terminal tail with other activin receptor-like kinase proteins. ALK-1 is mainly expressed in endothelial cells regulating proliferation and migration in vitro and angiogenesis in vivo. Mutations in ALK-1 as well as in endoglin are associated with hereditary hemorrhagic telangiectasia (HHT), suggesting ALK-1 plays a critical role for in the control of blood vessel development or repair.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.