


概述 (Summary)

英文全称	BTN1A1/Butyrophilin Subfamily 1 Member A1
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Butyrophilin Subfamily 1 Member A1 is produced by our Mammalian expression system and the target gene encoding Ala27-Trp247 is expressed with a 6His tag at the C-terminus.
Accession #	Q62556
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Mouse
预测分子量 (Predicted MW)	25.4 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse BTN1A1 (C-6His)
Catalog #: PHM0175



背景 (Background)

分子别名 (Alternative Names)

Butyrophilin subfamily 1 member A1; BTN; BTN1A1; butyrophilin; Btn1a1

背景介绍 (References)

Mouse Butyrophilin subfamily 1 member A1(BTN1A1) is a type I transmembrane glycoprotein which is a member of the Ig superfamily. The BTN1A1 ECD displays two predicted IgV and IgC domains as do B7 and Skint proteins which interact with other Ig superfamily members. BTN1A1 binds to xanthine oxidoreductase (XOR). This interaction stabilizes the association of XOR with the milk fat globule membrane and appears to be essential in the control of milk fat globule secretion. In vitro, BTN1A1 inhibits the proliferation of CD4 and CD8 T-cells activated by anti-CD3 antibodies, T-cell metabolism and IL-2 and IFN- γ secretion. Furthermore, in vivo, BTN1A1 has a protective effect against the development of experimental autoimmune encephalomyelitis (EAE). Because butyrophilins are structurally related to B7 proteins and are functionally implicated in immune regulation, they may represent an emerging family of costimulatory/inhibitory molecules.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.